DOI: 10.1111/1471-0528.18006 www.bjog.org



Abstracts

245

Estimated fetal weight versus birthweight discrepancy in large babies suggests slowing of growth before birth

Oliver Hugh, Jason Gardosi

Perinatal Institute, Birmingham, United Kingdom

Objectives: Ultrasound-estimated fetal weight (EFW) often overestimates the weight centile of the baby at birth, resulting in high rates of false positive assessments of large-for-gestational-age (LGA) fetuses.¹ We investigated whether the discrepancy could also be due to reduced growth velocity before birth.

Methods: We examined a routinely recorded database of 25,230 pregnancies that had two or more third-trimester scans for various indications. We selected a cohort of all cases where the last two EFWs were above the ninetieth customised centile (n= 2,501). We then compared this with the centile at birth, at the following intervals from last scan: <3, 4–6, 7–13, 14–20 and >21 days.

Results: The average gestational age (weeks + days) was 33+5 and 36+4 at the last two scans, and 38+6 at birth (interval 16 days from last scan). The average EFW centiles at the two scans were 97.1 and 97.8, respectively. The average centile at birth was 91.8, which included 46% of cases that were not LGA (<90th centile). Births within three days of the last scan showed a systematic error of 1.9 centile only. The centile drop at birth increased with the interval between last scan and birth (p < 0.01, Spearman Rank test) (see Figure). The pattern was similar for spontaneous-onset labours.

Conclusions: Nearly half of fetuses found to be LGA at last scan are not >90th centile at birth. The reduction of size appears to be due to a gradual slowing of growth at term.

Reference:

1. Ewington L, Hugh O, Butler E, et al. Accuracy of antenatal ultrasound in predicting large for gestational age babies. Submitted abstract, BMFMS 2024.



FIGURE 1